

UNEXPECTED BIASES IN PRIME FACTORIZATIONS AND LIOUVILLE FUNCTIONS FOR ARITHMETIC PROGRESSIONS

SNEHAL M. SHEKATKAR AND TIAN AN WONG

ABSTRACT. We introduce a refinement of the classical Liouville function to primes in arithmetic progressions. Using this, we discover new biases in the appearances of primes in a given arithmetic progression in prime factorizations of integers. For example, we observe that the primes of the form $4k+1$ tend to appear an even number of times in the prime factorization of a given integer, more so than for primes of the form $4k+3$. We also consider variants of Pólya's conjecture, supported by extensive numerical evidence, and its relation to other well-known conjectures.

CONTENTS

1. Introduction	1
2. First estimates	7
3. Parity	10
4. Distribution	20
5. Numerical tests	22
References	24

1. INTRODUCTION

1.1. The Liouville function. The classical Liouville function is the completely multiplicative function defined by $\lambda(p) = -1$ for any prime p . It can be expressed as

$$(1.1) \quad \lambda(n) = (-1)^{\Omega(n)}$$

where $\Omega(n)$ is the total number of prime factors of n . One sees that it is -1 if n has an odd number of prime factors, and 1 otherwise. By its relation to the Riemann zeta function

$$(1.2) \quad \sum_{n=1}^{\infty} \frac{\lambda(n)}{n^s} = \frac{\zeta(2s)}{\zeta(s)},$$

the Riemann hypothesis is known to be equivalent to the statement that

$$(1.3) \quad \sum_{n \leq x} \lambda(n) = O(x^{1/2+\epsilon})$$

Date: April 27, 2017.

2010 Mathematics Subject Classification. 11A51 and 11F68.

Key words and phrases. Liouville function, prime factorization, arithmetic progressions.

for any $\epsilon > 0$; whereas the prime number theorem is equivalent to the estimate $o(x)$. Indeed, the behaviour of the Liouville function, being a close relative of the more well-known Möbius function, is strongly connected to prime number theory. Also, we note that by the generalized Riemann hypothesis, one also expects (1.3) to hold for partial sums of $\lambda(n)$ restricted to arithmetic progressions.

In this paper, we introduce natural refinements of the Liouville function, which detect how primes in given arithmetic progressions appear in prime factorizations. Interestingly, we find that these functions behave in somewhat unexpected ways, which is in turn related to certain subtleties of the original Liouville function. We now describe this briefly.

Define $\Omega(n; q, a)$ to be the total number of prime factors of n congruent to $a \pmod q$, and

$$(1.4) \quad \lambda(n; q, a) = (-1)^{\Omega(n; q, a)}$$

to be the completely multiplicative function that is -1 if n has an odd number of prime factors congruent to $a \pmod q$, and 1 otherwise. They are related to the classical functions by

$$(1.5) \quad \lambda(n) = \prod_{a=0}^{q-1} \lambda(n; q, a), \quad \Omega(n) = \sum_{a=0}^{q-1} \Omega(n; q, a).$$

Using this we study the asymptotic behaviour instead of

$$(1.6) \quad L(x; q, a) = \sum_{n \leq x} \lambda(n; q, a),$$

hence the distribution of the values of $\lambda(n; q, a)$. Also, we will be interested in r -fold products of $\lambda(n; q, a)$,

$$(1.7) \quad \lambda(n; q, a_1, \dots, a_r) = \prod_{i=1}^r \lambda(n; q, a_i)$$

where the a_i are residue classes mod q , with $1 \leq r \leq q$, and define $\Omega(n; q, a_1, \dots, a_r)$ and $L(x; q, a_1, \dots, a_r)$ analogously.

1.2. Prime factorizations. Given a prime number p , we will call the *parity* of p in an integer n to be even or odd, according to the exponent of p in the prime factorization of n . This includes the case where p is prime to n , in which case its exponent is zero and therefore having even parity.


Landau showed that the number of $n \leq x$ containing an even (resp. odd) number of prime factors both tend to

$$(1.8) \quad \frac{1}{2}x + O(xe^{c\sqrt{\log x}})$$

with x tending to infinity, and c some positive constant. Soon after, Pólya asked whether the sum

$$(1.9) \quad \sum_{n \leq x} \lambda(n) \leq 0$$

for all $x > 2$; this was shown to be false by Haselgrove [H] using the zeroes of $\zeta(s)$, and that in fact the sum must change sign infinitely often. Indeed, the first sign change was later computed to be around 9×10^8 . A similar was problem posed by Turán on the positivity of partial sums of $\lambda(n)/n$, which was also shown to be false,

FIGURE 1. $L(x; 2, 2)$ and $L(x; 2, 1)$.

with the first sign change taking place around 7×10^{13} . (See [MRT] for a discussion of these problems.)

On the other hand, by the equidistribution of primes in arithmetic progressions, one might guess that the number of $n \leq x$ containing an even (resp. odd) number of prime factors $p \equiv a \pmod q$, for a fixed arithmetic progression would be evenly distributed over residue classes coprime to q . By our analysis of $\lambda(n; q, a)$, we find that this seems not to be the case. We view this as a different kind of ‘unexpected bias’, in contrast to [LOS], who uncovered biases in consecutive primes, and show its connection to the Hardy-Littlewood k -tuples conjecture.


For example, if we consider the parity of 2 and the parity of the odd primes separately, we find the behaviour of the partial sums as in Figure 1 above. Indeed, we prove that the sum $L(x; 2, 1)$ is $o(x)$, and $O(x^{1/2+\epsilon})$ on the Riemann hypothesis; whereas $L(x; 2, 2) \geq 0$ and tends to $\frac{1}{3}x$ unconditionally. Numerically, we find that $L(x; 2, 1) \leq 0$ at least up to $x \leq 10^9$. Interestingly, in spite of the expected square-root cancellation of $L(x; 2, 1)$, the graph suggests that $\lambda(x; 2, 1)$ favors -1 more than 1 . That is, separating the primes dividing the modulus appears to ‘tame’ the randomness of $\lambda(n)$.

More generally, we have the following theorem.

Theorem 1.1. *Given any $q \geq 2$, let $a_1, \dots, a_{\varphi(q)}$ be the residue classes mod q such $(a_i, q) = 1$, and $b_1, \dots, b_{q-\varphi(q)}$ the remaining residues classes. Then*

$$(1.10) \quad \sum_{n \leq x} \lambda(n; q, a_1, \dots, a_{\varphi(q)}) = o(x),$$

for $x \geq 1$. Assuming the Riemann hypothesis, it is in fact $O(x^{1/2+\epsilon})$ for all $\epsilon > 0$.

FIGURE 2. $L(x; 4, 3)$ and $L(x; 4, 1)$.

On the other hand,


$$(1.11) \quad \sum_{n \leq x} \lambda(n; q, b_1, \dots, b_k) = \left(\prod_{i=1}^k \prod_{p|b_i} \frac{p-1}{p+1} \right) x + o(x)$$

for any $1 \leq k \leq q - \varphi(q)$.

In fact, it is straightforward to show that the estimate $O(x^{1/2+\epsilon})$ is equivalent to the Riemann hypothesis, as in the classical Liouville function.

Having separated the odd primes, the natural thing to do next is to consider primes mod 4. Surprisingly, we observe the behaviour as in Figure 2 above. In this case, the arithmetic function $\lambda(n; 4, 3)$ resembles the non-principal Dirichlet character mod 4, and its partial sums are shown to be *positive*. On the other hand, the behaviour of $\lambda(n; 4, 1)$ turns out to be related to the classical $\lambda(n)$ restricted to arithmetic progressions mod 4. Numerically, we checked that the $L(x; 4, 1) \geq 0$ up to $x \leq 10^{11}$, but for arbitrary x we were only able to prove the weak estimate in Lemma 3.11.

For general moduli, taking $r = \varphi(q)/2$, for certain choices of a_1, \dots, a_r coprime to q , the function $\lambda(n; q, a_1, \dots, a_r)$ resembles a Dirichlet character, and in this sense is ‘character-like,’ in the sense of [BCC]. In this case, we can predict the behaviour of the function and its ‘complement’, that is, when we are in the analogous setting to $\lambda(n; 4, 1)$ and $\lambda(n; 4, 3)$.

FIGURE 3. Combinations of $L(x; 5, *, *)$.

Theorem 1.2. *Let a_1, \dots, a_{q-r} and b_1, \dots, b_r be as above. Then*

$$(1.12) \quad \sum_{n \leq x} \lambda(n; q, a_1, \dots, a_{q-r}) = O(x^{1/2+\epsilon})$$

is equivalent to the Riemann hypothesis for $L(s, \chi_q)$; unconditionally, it is $o(x)$. On the other hand,

$$(1.13) \quad \sum_{n \leq x} \lambda(n; q, b_1, \dots, b_r) = O(\log x).$$

if there is only one non-principal real Dirichlet character mod q , if there is more than one then we have only $o(x^{1-\delta})$ for some $\delta > 0$.

Moreover, we show that when $r \neq \varphi(q)/2$, and $(a_i, q) = 1$, the behaviour of $\lambda(n; q, a_1, \dots, a_r)$ is determined. Otherwise, the behaviour of $\lambda(n; q, a_1, \dots, a_r)$ seems more difficult to describe precisely, and in this case it is interesting to ask the same question as Pólya did for $\lambda(n)$. For example, with modulus 5 we observe as in Figure 3, that except for the character-like function and its complement, the partial sums tend to fluctuate with a positive bias, except for $\lambda(n; 5, 1, 2)$, which already changes sign for small x . The remaining three remain positive up to $x \leq 10^7$, which leads us to ask whether they eventually change sign. (See Problem 5.1.)

Theorem 1.3. *Let a_1, \dots, a_r be residue classes mod q , coprime to q . Then for $r \neq \varphi(q)/2$,*

$$(1.14) \quad \sum_{n \leq x} \lambda(n; q, a_1, \dots, a_r) = b_0 x (\log x)^{2r/\varphi(q)-2} + O(x (\log x)^{2r/\varphi(q)-3})$$

where b_0 is an explicit constant such that $b_0 > 0$ if $2r < \varphi(q)$ and $b_0 < 0$ if $2r > \varphi(q)$.

If $r = \varphi(q)/2$ and $\lambda(n; q, a_1, \dots, a_r)$ is not character like, we have again

$$(1.15) \quad \sum_{n \leq x} \lambda(n; q, a_1, \dots, a_{\varphi(q)}) = o(x),$$

for $x \geq 1$. Assuming the Riemann hypothesis, it is in fact $O(x^{1/2+\epsilon})$ for all $\epsilon > 0$.

The most intriguing aspect of our new family of Liouville-type functions, in light of the conjectures of Pólya, Turán, and even Mertens, is distinguishing when the partial sums of $\lambda(n; q, a_1, \dots, a_r)$ have any sign changes at all, or if a sign change occurs, then there must be infinitely many sign changes must follow. Our theorems only answer this question partially, and we speculate on this in Problem 5.1.

We also prove some results on distribution of total number of primes in arithmetic progression, though the analogues $\Omega(n; q, a)$ and $\omega(n; q, a)$ are more well-behaved, though we still observe some slight discrepancy in the implied constants in the growth of the partial sums, with respect to the residue class. For example, we show that

$$(1.16) \quad \sum_{n \leq x} \omega(n; q, a) = \frac{1}{\varphi(q)} x \log \log x + O\left(\frac{1}{\phi(q)} \frac{x}{\log x}\right)$$

(see Proposition 4.1), and that $\omega(n; q, a)$ is distributed normally, as an application of the Erdős-Kac theorem.

Note that we have not considered the ‘mixed’ case, where $\lambda(n; a_1, \dots, a_r)$ contains both residue classes that are and are not coprime to q . Numerical experiments seem to suggest that they do affect the behaviour in small but observable ways, in particular, we observe that the adding several residue classes may cause a sum to fluctuate more. See Section 5 for a discussion and an example.

Remark 1.4. We mention the recent work of [BCC], in which the authors consider any subset A of prime numbers, and define $\Omega_A(n)$ to be the number of prime factors of n contained in A , counted with multiplicity. They then define a Liouville function for A to be

$$(1.17) \quad \lambda_A(n) = (-1)^{\Omega_A(n)},$$

taking value -1 at primes in A and 1 at primes not in A , and show, for example, that $\lambda_A(n)$ is not eventually periodic in n . Our functions can be viewed as particular cases of $\lambda_A(n)$ where A is a set of primes in a given arithmetic progression.

1.3. This paper. This paper is organized as follows. In Section 2 we develop basic properties of the Liouville function for arithmetic progressions. In Section 3, we study the parity of the number of primes in arithmetic progressions appearing in a given prime factorization, while in Section 4 we study the distribution of the total number of such primes.

In Section 5, we discuss numerical experiments and the observed behaviour of these Liouville functions. Based on these, we pose several problems in Problem 5.1 in comparison to the original problem of Pólya. We note that the techniques used in this paper are mainly complex analytic, and the estimates obtained are not necessarily optimal.

Notation. Throughout the paper we fix the following conventions: ϵ will denote a positive real number and c an absolute constant, which will vary depending on

the context. We also denote by $[x]$ the floor function, or the greatest integer less than x .

Acknowledgments. The second author would like to thank Neha Prabhu for helpful discussions.

2. FIRST ESTIMATES

2.1. Basic properties. We develop some basic properties of the Liouville function for arithmetic progressions, analogous to the classical results. Using this we prove a basic estimate for the distribution of $\lambda(n; q, a)$. Most of the statements in this section will be proven for $\lambda(n; q, a)$, and we leave to the reader the analogous statements for products $\lambda(n; q, a_1, \dots, a_r)$.

We begin with a broad description.

Lemma 2.1. *$\lambda(n; q, a)$ is aperiodic and unbounded.*

Proof. The first statement follows as a special case of [BCC]. The second follows, for example, from the Erdős discrepancy problem,¹ but we will also prove this more directly below. \square

Recall that the classical Liouville function satisfies the identity

$$(2.1) \quad \sum_{d|n} \lambda(d) = \begin{cases} 1 & \text{if } n \text{ is a perfect square,} \\ 0 & \text{otherwise.} \end{cases}$$

The following proposition gives the analogue of this identity.

Lemma 2.2. *Let $n = n_1 n_2$ where n_1 is not divisible by any $p \equiv a \pmod{q}$. Then we have*

$$(2.2) \quad S(n; q, a) := \sum_{d|n} \lambda(d; q, a) = \begin{cases} \tau(n_1) & \text{if } n_2 \text{ is a perfect square,} \\ 0 & \text{otherwise.} \end{cases}$$

where $\tau(n)$ is the divisor function.

Proof. We will proceed by induction on k , writing $n_2 = p_1^{r_1} \dots p_k^{r_k}$. First suppose $k = 0$, then by definition $\lambda(n_1; q, a) = 1$, and therefore $S(n_1; q, a) = \tau(n_1)$. On the other hand, if $k = 1$ and r_1 is odd, then $n = n_1 p_1^{r_1}$ and

$$(2.3) \quad S(n; q, a) = \sum_{d|a} \sum_{i=0}^{r_1} \lambda(n_1 p_1^i; q, a) = \sum_{d|a} \lambda(n_1) \sum_{i=0}^{r_1} (-1)^i,$$

and the inner sum vanishes since r_1 is odd.

Now suppose that (2.2) holds for $k - 1$, and define A such that $n = A p_k^{r_k}$. Then

$$(2.4) \quad S(n; q, a) = \sum_{d|A} \sum_{i=0}^{r_k} \lambda(A p_k^i; q, a),$$

which, by multiplicativity, is

$$(2.5) \quad \left(\sum_{d|A} \lambda(d; q, a) \right) \left(\sum_{i=0}^{r_k} \lambda(p_k^i; q, a) \right) = S(A; q, a) \sum_{i=0}^{r_k} (-1)^i,$$

¹It may seem like big hammer to invoke, but it is worth noting that amongst our $\lambda(n; q, a)$ are ‘character-like’ multiplicative functions considered in [BCC], whose $O(\log x)$ growth constitute near misses to the problem.

since $\lambda(p_k^i; q, a) = (-1)^i$. Now we see that if n_2 is a perfect square, the sum is equal to 1 and $S(A; q, a) = \tau(A)$ by hypothesis; whereas if n_2 is not a perfect square, either r_k is odd and the sum vanishes, or r_k is even and $r_{k'}$ is odd for some $k' < k$, in which case $S(A; q, a) = 0$ by hypothesis. \square

From this we have a crude first estimate:

Corollary 2.3. $L(x; q, a) = O(x \log x)$ for any a, q , and $x > 1$.

Proof. This follows from the Möbius inversion formula

$$(2.6) \quad \lambda(n; q, a) = \sum_{d|n} \mu\left(\frac{n}{d}\right) S(d; q, a),$$

and by applying the trivial bounds on $\mu(n)$ and $S(n; q, a)$. \square

We will now do slightly better, at the expense of less elementary methods. We say a subset A of primes is said to have sifting density κ , if

$$(2.7) \quad \sum_{p \in A: p \leq x} \frac{\log p}{p} = \kappa \log x + O(1).$$

where $0 \leq \kappa \leq 1$. In particular, we will take A to be the set of primes congruent to certain $a \pmod q$. For example, we have $\kappa = 0$ if $(a, q) > 1$, and for $(a, q) = 1$ with q odd, we have $0 < \kappa \leq \frac{1}{2}$ with equality only when $\varphi(q) = 2$.

Proposition 2.4. *We have for $\kappa < \frac{1}{2}$, and $(a, q) = 1$,*

$$(2.8) \quad \sum_{n \leq x} \lambda(n; q, a) = (1 + o(1)) \frac{C_\kappa x}{(\log x)^{2\kappa}}$$

where $C_\kappa > 0$ is an explicit constant depending on a, q and κ , and for $\kappa \geq \frac{1}{2}$,

$$(2.9) \quad \sum_{n \leq x} \lambda(n; q, a) = o(x).$$

Proof. This follows from [BCC, Theorem 5], as an application of the Liouville function for A , choosing A to be a set of primes in arithmetic progression. \square

Remark 2.5. More generally, if we take A to consist of several residue classes $a_i \pmod q$, then κ will also vary accordingly according to the number of residue classes prime to q that are taken. In this case, we may replace the sum over $\lambda(n; q, a)$ by $\lambda(n; q, a_1, \dots, a_r)$ to obtain similar estimates.

We also record the following estimates for the classical Liouville function as a benchmark.

Lemma 2.6. *We have*

$$(2.10) \quad \sum_{n \leq x} \lambda(n) = O(xe^{-c_1 \sqrt{\log x}}).$$

and

$$(2.11) \quad \sum_{\substack{n \leq x \\ n \in P}} \lambda(n) = -c_2 x (\log x)^{-r/\phi(q)-1} + O(x (\log x)^{r/\phi(q)-2})$$

where P is a set of r residue classes coprime to $q > 2$ and $c_1, c_2 > 0$.

Proof. The first is well known. The second follows by the relation of $\lambda(n)$ to the Möbius function

$$(2.12) \quad \sum_{n \leq x} \lambda(n) = \sum_{n \leq \sqrt{x}} \sum_{m \leq x/m^2} \mu(m)$$

and the fact that

$$(2.13) \quad \sum_{\substack{n \leq x \\ n \in P}} \mu(n) = -c_2 \frac{x}{(\log x)^{r/\phi(q)+1}} + O(x(\log x)^{r/\phi(q)-2}).$$

from [C, Theorem 2]. \square

In particular, we observe that when $\lambda(n)$ is restricted to arithmetic progressions (containing infinitely many primes), its partial sums tend to be negative. One can also show that its limiting distribution is negative using its relation to Lambert series.

2.2. Complete multiplicativity. We now use some general facts about completely multiplicative functions to show what one might expect to hold in our case. Here we have not strived to provide the best bounds, but rather those simplest to state to illustrate the general picture.

The first shows the values of $\lambda(n)$ restricted to residue classes is equidistributed.

Proposition 2.7. *Let f be a completely multiplicative function, $A, Q \geq 1$, $x > Q$ and $(a, q) = 1$. Then*

$$(2.14) \quad \sum_{\substack{n \leq x \\ n \equiv a(q)}} f(n) - \frac{1}{\phi(q)} \sum_{\substack{n \leq x \\ (n, q) = 1}} f(n) = O\left(\frac{x}{q\sqrt{\log A}}\right)$$

for all $q > Q$, except possibly for multiples of one of at most two exceptional moduli.

Proof. See [AGS]. \square

One may also consider a variant of the Chowla conjecture for the Liouville function: Fix a, q relatively prime. Given distinct integers h_1, \dots, h_k , fix a sequence of signs $\epsilon_j = \pm 1$ for $1 \leq j \leq k$. Then one would like to know whether

$$(2.15) \quad \sum_{n \leq x} \prod_{i=1}^k \lambda(n + h_i; q, a) = o(x).$$

for all k . In particular, the number of $n \leq x$ such that $\lambda(n + j; q, a) = \epsilon_j$ for all $1 \leq j \leq k$ is

$$(2.16) \quad \left(\frac{1}{2^k} + o(1) \right) x.$$

Roughly, this tells us that $\lambda(n; q, a)$ takes the value 1 or -1 randomly.

Using known results, we have the following evidence towards the conjecture.

Proposition 2.8. *For every $h \geq 1$ there exists $\delta(h) > 0$ such that*

$$(2.17) \quad \frac{1}{x} \left| \sum_{n \leq x} \lambda(n; q, a) \lambda(n + 1; q, a) \right| \leq 1 - \delta(h)$$

for all x large enough. Similarly,

$$(2.18) \quad \frac{1}{x} \left| \sum_{n \leq x} \lambda(n; q, a) \lambda(n+1; q, a) \lambda(n+2; q, a) \right| \leq 1 - \delta(h)$$

Proof. This follows as a special case of [MR]. \square

Remark 2.9. We also note that as an application of [MR, Corollary 5] on sign changes of certain multiplicative functions, there exists a constant C such that every interval $[x, x + C\sqrt{x}]$ contains a number with an even number of prime factors in a fixed arithmetic progression $a \bmod q$, and another one with an odd number of such prime factors.

3. PARITY

We now turn to the average behaviour of our $\lambda(n; q, a)$. A refinement of Pólya's problem leads us to ask: consider the prime factorization of a composite number n . Do the primes in arithmetic progressions tend to appear an even or odd number of times? As described in the introduction, we show that one encounters surprising biases, namely, that the answer depends strongly on the arithmetic progression chosen.

3.1. Dirichlet series. Since $\lambda(n; q, a)$ is completely multiplicative, we can form the Dirichlet series generating function

$$(3.1) \quad D(s; q, a) := \sum_{n=1}^{\infty} \frac{\lambda(n; q, a)}{n^s} = \zeta(s) \prod_{p \equiv a(q)} \frac{1 - p^{-s}}{1 + p^{-s}}$$

using the Euler product in the case $(a, q) = 1$, and by the trivial bound converges absolutely for $\operatorname{Re}(s) > 1$. Taking the product over all such a , we obtain

$$(3.2) \quad \prod_{(a, q)=1} D(s; q, a) = \zeta(2s) \zeta(s)^{\varphi(q)-2} \prod_{p|q} \frac{1 + p^{-s}}{1 - p^{-s}}.$$

and we see that in the region $\operatorname{Re}(s)$, the expression has a pole of order $\varphi(q) - 2$ at $s = 1$ and a simple pole at $s = \frac{1}{2}$. Moreover, if we include residue classes a such that $(a, q) > 1$, in which this case $D(s; q, a)$ is equal to $\zeta(s)$ up a finite number of factors, we have

$$(3.3) \quad \prod_{a=0}^{q-1} D(s; q, a) = \zeta(2s) \zeta(s)^{q-2},$$

generalizing the classical formula.

Similarly, for products $\lambda(n; q, a_1, \dots, a_r)$ with $(a_i, q) = 1$ for each i , we have

$$(3.4) \quad D(s; q, a_1, \dots, a_r) = \zeta(s) \prod_{i=1}^r \prod_{p \equiv a_i(q)} \frac{1 - p^{-s}}{1 + p^{-s}}$$

and

$$(3.5) \quad D(s; q, a_1, \dots, a_r) D(s; q, a'_1, \dots, a'_{\varphi(q)-r}) = \zeta(2s) \prod_{p|q} \frac{1 + p^{-s}}{1 - p^{-s}}.$$

where $a'_1, \dots, a'_{\varphi(q)-r}$ are the remaining residue classes coprime to q .

Remark 3.1. We note in passing that by the non-negativity of the convolution $1 * \lambda(n; q, a_1, \dots, a_r)$, together with Dirichlet's hyperbola method, it is a pleasant exercise to show that any such Dirichlet series $D(s; q, a_1, \dots, a_r)$ is nonvanishing at $s = 1$, without appealing to the nonvanishing of $\zeta(s)$.

It is known that for $(a, q) = 1$, the partial product

$$(3.6) \quad F_a(s) = \prod_{p \equiv a(q)} \frac{1}{1 - p^{-s}}$$

converges absolutely for $\operatorname{Re}(s) > 1$, and has analytic continuation to $\operatorname{Re}(s) \geq 1 - C/\log t$ for $|t| < T$, and $T \geq 10$ [K, p.212]. It can be expressed as

$$(3.7) \quad \prod_{p \equiv a(q)} \frac{1}{1 - p^{-s}} = \zeta(s)^{1/\varphi(q)} e^{G_a(s)},$$

where $G_a(s)$ is given by

$$(3.8) \quad \frac{1}{\varphi(q)} \sum_{\chi \neq \chi_0} \chi(\bar{a}) \left(\log L(s, \chi) + \sum_p \sum_{m \geq 1} \frac{\chi(p) - \chi^m(p)}{mp^{ms}} \right) + \frac{1}{\varphi(q)} \sum_{p|q} \log(1 - p^{-s}),$$

where $a\bar{a} \equiv 1 \pmod{q}$, and χ runs over Dirichlet characters mod q .

Even though $\lambda(n; q, a)$ is not eventually periodic, we are still relate it at times to Dirichlet characters by the following identity.

Proposition 3.2. *Let $q > 2$. Given any χ_q is a non-principal real Dirichlet character mod q , there is a combination of residue classes, say a_1, \dots, a_r such that*

$$(3.9) \quad D(s; q, a_1, \dots, a_r) = \prod_{p|q} \frac{1}{1 - p^{-s}} L(s, \chi_q).$$

Proof. Given χ_q , it is clear that we can choose a combination of residue classes a_1, \dots, a_r , coprime to q such that

$$(3.10) \quad \lambda(n; q, a_1, \dots, a_r) = \chi_q(n)$$

for any n in $(\mathbf{Z}/q\mathbf{Z})^\times$. Then it is clear that we may express the Dirichlet series as

$$(3.11) \quad D(s; q, a_1, \dots, a_r) = \prod_p (1 - \lambda(p; q, a_1, \dots, a_r) p^{-s})^{-1}$$

$$(3.12) \quad = \prod_{p|q} (1 - p^{-s})^{-1} \prod_{p \nmid q} (1 - \chi_{q,a}(p) p^{-s})^{-1},$$

and the result follows. \square

Finally, note that by the general theory of Dirichlet series we can relate it to the summatory function by

$$(3.13) \quad D(s; q, a) = s \int_1^\infty L(x; q, a) x^{-s-1} dx$$

for $\operatorname{Re}(s)$ large enough. We will use this expression implicitly throughout.

Remark 3.3. From the Möbius inversion formula it follows then that

$$(3.14) \quad \sum_{n=1}^\infty \frac{S(n; q, a)}{n^s} = D(s; q, a) \zeta(s) = \zeta(s)^2 \prod_{p \equiv a(q)} \frac{1 - p^{-s}}{1 + p^{-s}}$$

where the second equality follows from (3.1), but we will not use this relation in this paper.

3.2. The odd primes and 2. The first natural refinement is to ask what is the parity of (i) the odd primes and (ii) the prime 2 in prime factorizations. Indeed, we have:

Proposition 3.4.

$$(3.15) \quad \sum_{n \leq x} \lambda(n; 2, 1) = O(x^{1/2+\epsilon}),$$

for $x \geq 1$, assuming the Riemann hypothesis, for all $\epsilon > 0$. Otherwise, we have $o(x)$.² On the other hand,

$$(3.16) \quad \sum_{n \leq x} \lambda(n; 2, 2) = \frac{1}{3}x + o(x)$$

and is nonnegative for all $x \geq 1$.

Proof. We first treat the simpler case $L(n; 2, 2)$. The Dirichlet series

$$(3.17) \quad D(s; 2, 2) = \frac{1 - 2^{-s}}{1 + 2^{-s}} \zeta(s)$$

has meromorphic continuation to $\operatorname{Re}(s) \geq 1$ with only a simple pole at $s = 1$ with residue $\frac{1}{3}$, and is holomorphic for $\operatorname{Re}(s) > 1$, hence we have

$$(3.18) \quad \sum_{n \leq x} \lambda(n; 2, 2) = \frac{1}{3}x + o(x)$$

by the Ikehara-Wiener theorem.

Now, notice that $\lambda(n; 2, 2)$ is always $1, -1, 1$ when n is of the form $4k + 1, 4k + 2, 4k + 3$ respectively. Only if it is of the form $4k$ it can take 1 or -1 , in which case it is determined by the value $\lambda(k; 2, 2)$. Thus the first few summands of $L(x; 2, 2)$ are

$$(3.19) \quad 1 - 1 + 1 + \lambda(4; 2, 2) + 1 - 1 + 1 + \lambda(8; 2, 2) + \dots$$

and continuing thus, we conclude that for $L(x; 2, 2) \geq 0$ for all $x \geq 1$.

Next, we treat the case $L(x; 2, 1)$. Notice that

$$(3.20) \quad D(s; 2, 1) = \frac{1 + 2^{-s}}{1 - 2^{-s}} \frac{\zeta(2s)}{\zeta(s)}$$

where we recall from (1.2) that $\zeta(2s)/\zeta(s)$ is the Dirichlet series for the classical Liouville function. Since $\zeta(s)$ has a simple pole at $s = 1$ and is nonvanishing on $\operatorname{Re}(s) = 1$, it follows that $D(s; 2, 1)$ has analytic continuation to $\operatorname{Re}(s) \geq 1$, and $D(1; 2, 1) = 0$. Thus we have

$$(3.21) \quad \sum_{n \leq x} \lambda(n; 2, 1) = o(x)$$

unconditionally. As with the classical Liouville function, assuming the Riemann hypothesis, we see that $D(s; 2, 1)$ continues analytically to $\operatorname{Re}(s) > \frac{1}{2}$, and has a simple pole at $s = \frac{1}{2}$, thus we find that $L(x; 2, 1) = O(x^{\frac{1}{2}+\epsilon})$. \square

²As in Figure 1, from numerical evidence we expect it to be in fact always negative for $x > 2$.

By a similar argument, we observe the following behaviour for general arithmetic progressions:

Theorem 3.5. *Given any $q \geq 2$, let $a_1, \dots, a_{\varphi(q)}$ be the residue classes mod q such $(a_i, q) = 1$, and $b_1, \dots, b_{q-\varphi(q)}$ remaining the residues classes. Then*

$$(3.22) \quad \sum_{n \leq x} \lambda(n; q, a_1, \dots, a_{\varphi(q)}) = O(x^{1/2+\epsilon}),$$

for $x \geq 1$, assuming the Riemann hypothesis, for all $\epsilon > 0$. Otherwise, we have $o(x)$.

On the other hand,

$$(3.23) \quad \sum_{n \leq x} \lambda(n; q, b_1, \dots, b_k) = \left(\prod_{i=1}^k \prod_{p|b_i} \frac{p-1}{p+1} \right) x + o(x)$$

for any $1 \leq k \leq q - \varphi(q)$.

Proof. Simply observe that the Dirichlet series in this setting can be expressed as

$$(3.24) \quad D(s; q, a_1, \dots, a_{\varphi(q)}) = \frac{\zeta(2s)}{\zeta(s)} \prod_{p|q} \frac{1+p^{-s}}{1-p^{-s}},$$

and

$$(3.25) \quad D(s; q, b_1, \dots, b_k) = \zeta(s) \prod_{i=1}^k \prod_{p|b_i} \frac{1-p^{-s}}{1+p^{-s}},$$

and argue as in Proposition 3.4. □

Remark 3.6. Qualitatively, we note that since $(p-1)/(p+1) < 1$ for any p , the growth rate becomes slower and slower as more p 's enter into the product. Numerically, we also observe that (3.22) appears to be non-positive for $x > 1$. See Section 5 for further discussion.

The following corollary is proved in the same manner for the classical Liouville function, after the method of Landau.

Corollary 3.7. *With assumptions as in Theorem 3.5,*

$$(3.26) \quad \sum_{n \leq x} \lambda(n; q, a_1, \dots, a_{\varphi(q)}) = o(x),$$

is equivalent to the prime number theorem.

By a similar reasoning, we shall see that one can also recover Dirichlet's theorem on primes in arithmetic progressions, but only in cases where there are no complex (i.e., non-real) Dirichlet characters mod q .

3.3. A Chebyshev-type bias. We show a result using the properties developed above, which can be interpreted as: the number of prime factors of the form $4k+1$ and $4k+3$ both tend to appear an even number of times, but the former having a much stronger bias. We first require the following formula.

Lemma 3.8. *Define the characteristic function*

$$(3.27) \quad c(x, k) = \begin{cases} 1 & 2k \leq x < 3k \pmod{4k} \\ 0 & \text{otherwise} \end{cases}$$

for any $x > 0$. Then

$$(3.28) \quad \sum_{n \leq x} \lambda(n; 4, 3) = \sum_{k=1}^{\infty} c(x, 2^k),$$

the with finitely many terms on the right-hand side being nonzero.

Proof. To prove the formula, we repeatedly apply the elementary fact that if $n \equiv 1 \pmod{4}$ (resp. $3 \pmod{4}$), then the prime factors of n of the form $3 \pmod{4}$ appear an even (resp. odd) number of times.

First, observe that $\lambda(n; 3, 4)$ is 1 or -1 if n is 1 or $3 \pmod{4}$, thus the sum

$$(3.29) \quad \sum_{\substack{n \leq x \\ n \equiv 1(2)}} \lambda(n; 4, 3)$$

is equal to $c(n, 2)$. Then we move on to the even numbers, which, written as $2m, 2(m+1)$ and using $\lambda(2; 4, 3) = 1$, gives again the pattern 1 and -1 depending on whether m is 1 or $3 \pmod{4}$. The even numbers of the form 4 and $6 \pmod{8}$ contribute the term $c(n, 2^2)$.

Repeating this process we obtain the terms $c(n, 2^k)$ for all k , but certainly for k large enough this procedure will cover all $n \leq x$, so only finitely many terms will be nonzero. \square

Proposition 3.9. *Let χ_4 be the non-principal Dirichlet character mod 4. Then*

$$(3.30) \quad \sum_{n \leq x} \lambda(n; 4, 1) = O(x^{1/2+\epsilon})$$

assuming the Riemann hypothesis for $L(s, \chi_4)$, otherwise we have $o(x)$ unconditionally. On the other hand,

$$(3.31) \quad \sum_{n \leq x} \lambda(n; 4, 3) = o(x^\epsilon)$$

for any $\epsilon > 0$ and is nonnegative for $x \geq 1$.

Proof. We first prove the latter statement. The Dirichlet series of $\lambda(n; 4, 3)$ can be written as

$$(3.32) \quad D(s; 4, 3) = \frac{1}{1 - 2^{-s}} L(s, \chi_4),$$

thus we see that $D(s; 4, 3)$ has analytic continuation to the half-plane $\operatorname{Re}(s) > 0$. Moreover, χ_4 being an even Dirichlet character one knows that $L(s, \chi_4)$ vanishes at $s = 0$, whereas the denominator $1 - 2^{-s}$ has a pole at $s = 0$.

From the explicit formula in Lemma 3.8 we also see immediately that $L(x; 4, 3)$ is nonnegative, and given any $C > 0$ we can find x large enough so that $L(x; 4, 3) > C$. Hence $D(s; 4, 3)$ has a simple pole at $s = 0$, and finally we conclude that

$$(3.33) \quad \sum_{n \leq x} \lambda(n; 4, 3) = o(x^\epsilon)$$

for any $\epsilon > 0$.

On the other hand, from (3.2) we have

$$(3.34) \quad D(s; 4, 1)D(s; 4, 3) = \zeta(2s) \frac{1 + 2^{-s}}{1 - 2^{-s}},$$

which by (3.32) is

$$(3.35) \quad D(s; 4, 1) = \frac{\zeta(2s)}{L(s, \chi_4)} (1 + 2^{-s})$$

Comparing both sides, we observe that $D(s; 4, 1)$ is analytic in the region $\operatorname{Re}(s) \geq 1$, giving $o(x)$ unconditionally by analytic continuation of $L(s, \chi_4)$. Moreover, assuming the Riemann hypothesis for $L(s, \chi_4)$, we see that $D(s; 4, 1)$ in fact converges absolutely in $\operatorname{Re}(s) > \frac{1}{2}$, with only a simple pole at $s = \frac{1}{2}$, so that

$$(3.36) \quad \sum_{n \leq x} \lambda(n; 4, 1) = o(x^\epsilon).$$

since $L(\frac{1}{2}, \chi_4) > 0$. □

Remark 3.10. Indeed, we can numerically observe that the growth of $L(x; 4, 1)$ is extremely slow: for $x \leq 10^7$ the maximum value attained is 14, while for $x \leq 10^9$ the maximum value is 29. Also, numerical experiments support the expectation that

$$(3.37) \quad \sum_{n \leq x} \lambda(n; 4, 1) \sim 2.557x^{1/2}$$

using the approximation $L(s, \chi) = 0.6677\dots$ at $s = \frac{1}{2}$. Unfortunately, we have only the poor unconditional estimate below.

Lemma 3.11. *Let χ'_{12}, χ''_{12} be the Dirichlet characters mod 12 such that $\chi(5) \neq \chi(11)$. Then*

$$(3.38) \quad \sum_{n \leq x} \lambda(n; 4, 1) = Cx^{1/2} - \frac{1}{2} \sum_{a \leq x} (\chi'_{12} - \chi''_{12})(a) \sum_{x/(a+2) < b \leq x/a} \lambda_4(b) + O(1)$$

where

$$(3.39) \quad C = \left(1 + \frac{1}{\sqrt{3}}\right) \sum_{k=0}^{\log_2 x} \frac{1}{2^{k/2+1}} = 2.6927\dots$$

Proof. We make the following reduction: since $\lambda(2n; 4, 1) = \lambda(n; 4, 1)$, we have a decomposition

$$(3.40) \quad \sum_{n \leq x} \lambda(n; 4, 1) = \sum_k \sum_{\substack{n \leq x/2^k \\ (n, 2)=1}} \lambda(n; 4, 1)$$

where the outer sums are taken over integers $0 \leq k \leq \log_2 x$. Thus it will suffice to estimate the inner sum. To ease notation, we will denote by $\lambda_4(n) = \lambda(n; 4, 1)$.

We begin with the observation that for n odd,

$$(3.41) \quad \chi_4 * \lambda_4 = \chi_4 * \chi_4 \lambda = 1_{n^2}$$

where $\lambda(n)$ is the classical Liouville function, and χ_4 is the Dirichlet character with sign pattern 1, -1 on residue classes 1 and 3 mod 4. Summing over up to x ,

$$(3.42) \quad \sum_{n \leq x} (\chi_4 * \lambda_4)(n) = \sum_{\substack{n \leq \sqrt{x} \\ n \equiv 1(2)}} 1 = \frac{1}{2} [\sqrt{x}] + O(1)$$

where the $O(1)$ is only due the ratio

$$(3.43) \quad \frac{\#\{n \text{ odd} \leq \sqrt{x}\}}{\#\{n \leq \sqrt{x}\}} \sim \frac{1}{2} \pm \frac{1}{\sqrt{x}}.$$

We may also absorb the error $[\sqrt{x}] = \sqrt{x} + O(1)$, but we maintain the expression for exactness. Expanding the convolution and using the cancellation in $\chi_4(n)$, we write the left hand side of (3.42) as

$$(3.44) \quad \sum_{a \leq x} \chi_4(a) \sum_{b \leq x/a} \lambda_4(b)$$

$$(3.45) \quad = \sum_{\substack{a \leq x \\ a \equiv 1(4)}} \sum_{x/(a+2) < b \leq x/a} \lambda_4(b)$$

$$(3.46) \quad = \sum_{n \leq x} \lambda_4(n) - \sum_{\substack{a \leq x \\ a \equiv 3(4)}} \sum_{x/(a+2) < b \leq x/a} \lambda_4(b)$$

with a, b both odd. This is the first step.

Second, replace x with $x/3$ in (3.46), so that

$$(3.47) \quad \sum_{\substack{a \leq x/3 \\ a \equiv 1(4)}} \sum_{x/3(a+2) < b \leq x/3a} \lambda_4(b) = \frac{1}{2} \left[\sqrt{\frac{x}{3}} \right] + O(1).$$

Adding this to the previous sum (3.46), we see that this completes the sum further, except for the intervals $[a, a+2]$ which are counted twice and $[a+6, a+8]$ not at all, where $a \equiv 5 \pmod{12}$. In fact, this is perfectly described by the Dirichlet character mod 12 = 3×4 , that takes the values 1, 1, -1, -1 on residue classes 1, 5, 7, and 11 mod 12, which we denote χ'_{12} . That is,

$$(3.48) \quad \sum_{n \leq x} \lambda_4(n) + \sum_{\substack{a \leq x \\ a \equiv 5(12)}} \sum_{x/(a+2) < b \leq x/a} \lambda_4(b) - \sum_{\substack{a \leq x \\ a \equiv 11(12)}} \sum_{x/(a+2) < b \leq x/a} \lambda_4(b)$$

simplifies to

$$(3.49) \quad \sum_{n \leq x} (\chi'_{12} * \lambda_4)(n) = \sum_{a \leq x} \chi'_{12}(a) \sum_{b \leq x/a} \lambda_4(b) = \frac{1}{2} [\sqrt{x}] + \frac{1}{2} \left[\sqrt{\frac{x}{3}} \right] + O(1),$$

the sum taken still over odd integers. Note that χ_4 restricted to $(\mathbf{Z}/12\mathbf{Z})^\times$ is equal to χ'_{12} have the same sign pattern. Apply Fourier inversion to the last two terms in (3.48) to get

$$(3.50) \quad \frac{1}{4} \sum_{\chi} (\bar{\chi}(5) - \bar{\chi}(11)) \sum_{a \leq x} \chi(a) \sum_{x/(a+2) < b \leq x/a} \lambda_4(b)$$

There four Dirichlet characters mod 12, all real characters. There are two characters for which $\bar{\chi}(5) \neq \bar{\chi}(11)$, one of which is χ'_{12} and the other we denote χ''_{12} . Thus we have

$$(3.51) \quad \frac{1}{2} \sum_{a \leq x} (\chi'_{12} - \chi''_{12})(a) \sum_{x/(a+2) < b \leq x/a} \lambda_4(b)$$

Then rearranging and summing over k proves the claim. \square

Remark 3.12. By analyzing (3.51) further, we see that may further add correction terms

$$(3.52) \quad \frac{1}{2} \left([\sqrt{x}] + \left\lfloor \sqrt{\frac{x}{3}} \right\rfloor - \left\lfloor \sqrt{\frac{x}{5}} \right\rfloor + \left\lfloor \sqrt{\frac{x}{7}} \right\rfloor \right) + O(1)$$

leading to a better approximation $C' = 2.574 \dots$ but a more complicated error term. Unfortunately, this still does not give good enough control over the remainder, nor is there a clear pattern for continuing with the correction terms.

The proposition above holds more generally for any $q \geq 2$, by the same method of proof, using the following observation: Let $r = \varphi(q)/2$. Then there is exactly one combination of residue classes, say b_1, \dots, b_r such that

$$(3.53) \quad D(s; q, b_1, \dots, b_r) = \prod_{p|q} \frac{1}{1 - p^{-s}} L(s, \chi_q)$$

where χ_q is a non-principal real Dirichlet character mod q ; whereas

$$(3.54) \quad D(s; q, a_1, \dots, a_{q-r}) = \frac{\zeta(2s)}{L(s, \chi_q)} \prod_{p|q} (1 - p^{-s}),$$

where a_1, \dots, a_{q-r} are the remaining residue classes. We first observe that it converges absolutely at $s = 1$, which implies that $\lambda(n; q, a_1, \dots, a_{q-r})$ can have at most finitely many sign changes.

The following theorem is now immediate.

Theorem 3.13. *Let a_1, \dots, a_{q-r} and b_1, \dots, b_r be as above. Then*

$$(3.55) \quad \sum_{n \leq x} \lambda(n; q, a_1, \dots, a_{q-r}) = O(x^{1/2+\epsilon})$$

is equivalent to the Riemann hypothesis for $L(s, \chi_q)$, otherwise it is $o(x)$ unconditionally. On the other hand,

$$(3.56) \quad \sum_{n \leq x} \lambda(n; q, b_1, \dots, b_r) = O(\log x).$$

if there is only one non-principal real Dirichlet character mod q , otherwise we have only $o(x^\epsilon)$.

Proof. The estimate (3.55) follows the same argument as before, and the fact that $L(\frac{1}{2}, \chi_q) > 0$ for any non-principal real Dirichlet character. For (3.56), we use the relation between χ_q and the Kronecker symbol, which follows by a modest extension of [BCC, Corollary 6] to composite q . \square

Remark 3.14. We have not strived for the optimal unconditional bounds, that is, not assuming the Riemann hypothesis. The relevant estimates can certainly be improved, for example, using the zero-free regions for the associated Dirichlet L -functions and $\zeta(s)$.

Remark 3.15. In the case where $\lambda(n; q, a_1, \dots, a_{q-r})$ is the complement to a character-like function, we observe that when n is prime to some $p|q$, we have for $(n, q) = 1$,

$$(3.57) \quad \lambda(n; q, a_1, \dots, a_{q-r}) = \lambda(n) \chi_q(n),$$

where $\lambda(n)$ is the classical Liouville function. Then by (3.40) we can decompose

$$(3.58) \quad \sum_{n \leq x} \lambda(n; q, a_1, \dots, a_{q-r}) = \sum_{k=0}^{\lfloor \log_p x \rfloor} (k+1) \sum_{i=1}^r \chi_q(a_i) \sum_{\substack{x/p^{k+1} < n \leq x/p^k \\ n \equiv a_i(q)}} \lambda(n),$$

so that we are led to studying the partial sums of $\lambda(n)$ in arithmetic progressions. Using Fourier inversion and the convolution identity $\chi * \chi \lambda = 1 * \lambda = 1$, we can write the inner sum as

$$(3.59) \quad \sum_{\substack{y < n \leq z \\ n \equiv a_i(q)}} \lambda(n) = \frac{1}{\phi(q)} \sum_{\chi \neq \chi_0} \bar{\chi}(a_i) \frac{1}{2\pi i} \int_{b-iT}^{b+iT} \frac{\zeta(2s)}{L(s, \chi)} \frac{x^s}{s} ds + O(\log z).$$

for $y < n \leq z$. Then by a variation of [HS, Y], one can, assuming the GRH prove that

$$(3.60) \quad \lambda(n; q, a_1, \dots, a_{q-r}) = O(x^{1/2} \exp((\log x)^{1/2} (\log \log x)^c))$$

for some explicit constant $c > 0$.

3.4. General arithmetic progressions. Now we turn to general arithmetic progressions. We now restrict to residue classes a coprime to q , which is the most interesting case. We will also assume moreover that $\varphi(q) > 2$.

Proposition 3.16. *Let a_1, \dots, a_r be residue classes mod q , coprime to q , with $r \neq \varphi(q)/2$. Then*

$$(3.61) \quad \sum_{n \leq x} \lambda(n; q, a_1, \dots, a_r) = b_0 x (\log x)^{2r/\varphi(q)-2} + O(x (\log x)^{2r/\varphi(q)-3})$$

where b_0 is an explicit constant such that $b_0 > 0$ if $2r < \varphi(q)$ and $b_0 < 0$ if $2r > \varphi(q)$.

Proof. The proof is a slight modification of [K, Theorem 1]. We will give the main steps of the argument, leaving the details to the reader. To ease notation, let $D(s) = D(s; q, a_1, \dots, a_r)$. Let also b_1, \dots, b_t be the remaining $t = \varphi(q) - r$ coprime residue classes. We can express it as

$$(3.62) \quad D(s) = \frac{\zeta(2s)}{\zeta(s)} \prod_{i=1}^t \prod_{p \equiv b_i(q)} \frac{1 - p^{-2s}}{(1 - p^{-s})^2}$$

$$(3.63) \quad = \frac{\zeta(2s)}{\zeta(s)} \prod_{i=1}^t \frac{F_{b_i}(s)^2}{F_{b_i}(2s)}$$

$$(3.64) \quad = \zeta(s)^{2r/\varphi(q)-1} \zeta(2s)^{1-r/\varphi(q)} \prod_{i=1}^t \exp(2G_{b_i}(s) - G_{b_i}(2s))$$

by (3.7). Applying Perron's formula, we have

$$(3.65) \quad \sum_{n \leq x} \lambda(n; q, a_1, \dots, a_r) = \frac{1}{2\pi i} \int_{b-iT}^{b+iT} D(s) \frac{x^s}{s} ds + O\left(\frac{x}{T}\right)$$

for $b > 1$ and $x, T \geq 2$.

We may analytically continue $D(s)$ to the left of the line $\operatorname{Re}(s) = 1$, say to $\sigma \geq 1 - C/\log T$, so that the integral may be estimated by

$$(3.66) \quad \frac{1}{2\pi i} \int_{b-iT}^{b+iT} D(s) \frac{x^s}{s} ds = \frac{1}{2\pi i} \int_{\gamma} D(s) \frac{x^s}{s} ds + O\left(\frac{x}{T}\right)$$

where γ is a closed loop around $s = 1$, with radius taken to be less than $1 - C/\log T$. In fact, we will choose T such that $\log T = C(\log x)^{1/2}$.

Now if we define the function $H(s)$ by the equation

$$(3.67) \quad \frac{D(s)}{s} = (s-1)^{1-2r/\varphi(q)} H(s)$$

we can write

$$(3.68) \quad \frac{1}{2\pi i} \int_{\gamma} D(s) \frac{x^s}{s} ds = xI(x)$$

where

$$(3.69) \quad I(x) = \frac{1}{2\pi i} \int_{\gamma'} H(s+1) x^s s^{1-2r/\varphi(q)} ds$$

and γ' is the contour obtained by translating γ by $s \mapsto s+1$. Taking γ to have radius $1/N$, the integral can be written as (c.f. [K, p.212])

$$(3.70) \quad I(x) = \sum_{0 \leq j \leq N} B_j \frac{(\log x)^{r/\varphi(q)-j-2}}{\Gamma(2r/\varphi(q)-j-1)} + O\left((\log x)^{2r/\varphi(q)-N-2}\right)$$

where B_j are the coefficients of the Taylor series expansion of $H(s)$ at $s = 1$. In particular, $B_0 = H(1) > 0$. Set

$$(3.71) \quad b_j = B_j \frac{(\log x)^{r/\varphi(q)-j-2}}{\Gamma(2r/\varphi(q)-j-1)},$$

and notice that $\Gamma(2r/\varphi(q)-1)$ is positive or negative depending on whether $2r/\varphi(q)$ is lesser or greater than 1, and is singular at $2r = \varphi(q)$.


Finally, putting this together we have

$$(3.72) \quad \sum_{n \leq x} \lambda(n; q, a_1, \dots, a_r) = b_0 x (\log x)^{2r/\varphi(q)-2} + O(x (\log x)^{r/\varphi(q)-3})$$

as desired. \square

Remark 3.17. The above proposition is illustrated in the case $q = 5$ as in Figure 4. Indeed, from the graphs one would be led to ask if not only should the functions be asymptotically positive (resp. negative), but in fact positive (resp. negative) for all $x \geq 1$.

We now consider the most interesting case.

FIGURE 4. $L(x; 5, 1)$ and $L(x; 5, 1, 2, 3)$.

Proposition 3.18. *Let a_1, \dots, a_r be residue classes mod q , coprime to q , with $r = \varphi(q)/2$. Then for any $\lambda(n; q, a_1, \dots, a_r)$ that is neither character-like or the complement thereof, the sum*

$$(3.73) \quad \sum_{n \leq x} \lambda(n; q, a_1, \dots, a_r) = O(x^{1/2+\epsilon})$$

assuming the Generalized Riemann hypothesis.

Proof. The proof follows from a simple application of Ingham's analysis of $\lambda(n)$. Recall the Dirichlet series expression from (3.64), we have

$$(3.74) \quad \sum_{n \leq x} \lambda(n; q, a_1, \dots, a_r) = \zeta(2s)^{1/2} \prod_{i=1}^r \exp(2G_{a_i}(s) - G_{a_i}(2s))$$

where we have used $r = \varphi(q)/2$. □

4. DISTRIBUTION

In this section we provide a brief discussion on the distribution of the primes in arithmetic progressions in the number of prime factors.

4.1. The number of prime factors. Recall the functions $\omega(n)$ counting the number of distinct prime factors of n , and $\Omega(n)$ counting the total number of prime factors of n . We may express them as

$$(4.1) \quad \omega(n) = \sum_{p|n} 1, \quad \Omega(n) = \sum_{j=1}^{\infty} \sum_{p:p^j|n} 1.$$

As in (1.5) we may define the analogous functions $\omega(n; q, a)$ and $\Omega(n; q, a)$ counting only primes congruent to $a \pmod q$, so that

$$(4.2) \quad \omega(n) = \sum_{a=0}^{q-1} \omega(n; q, a), \quad \Omega(n) = \sum_{a=0}^{q-1} \Omega(n; q, a).$$

Proposition 4.1. *We have*

$$(4.3) \quad \sum_{n \leq x} \omega(n; q, a) = \frac{1}{\varphi(q)} x \log \log x + O\left(\frac{1}{\phi(q)} \frac{x}{\log x}\right)$$

Proof. Write

$$(4.4) \quad \sum_{n \leq x} \omega(n; q, a) = \sum_{n \leq x} \sum_{\substack{p|n \\ p \equiv a(q)}} 1 = \sum_{\substack{p \leq x \\ p \equiv a(q)}} \sum_{m \leq x/p} 1$$

which is

$$(4.5) \quad \sum_{\substack{p \leq x \\ p \equiv a(q)}} \frac{x}{p} + O\left(\sum_{\substack{p \leq x \\ p \equiv a(q)}} 1\right)$$

and by Dirichlet's theorem for primes in arithmetic progressions,

$$(4.6) \quad \sum_{\substack{p \leq x \\ p \equiv a(q)}} \frac{x}{p} + O\left(\frac{1}{\phi(q)} \frac{x}{\log x}\right).$$

Now, using Mertens' theorem for primes in arithmetic progressions for $(a, q) = 1$ we have

$$(4.7) \quad \sum_{\substack{p \leq x \\ p \equiv a(q)}} \frac{1}{p} = \frac{1}{\phi(q)} \log \log x + g(q, a) + o(1)$$

where $g(q, a)$ is an absolute constant. Applying this yields the proposition. \square

Remark 4.2. We note that while the main term in Prop 4.1 is independent of a , the error term does appear to depend on the choice of residue class a (see Figure 5 below), but we do not study this here.

We may also consider moments, such as

$$(4.8) \quad \sum_{n \leq x} (\omega(n; q, a) - \frac{1}{\phi(q)} \log \log x)^2 = \frac{x}{\phi(q)} \log \log x + O(x)$$


by expanding the square and applying simple estimates.

Moreover, since $\omega(n; q, a)$ is completely additive, we may apply the Erdős-Kac theorem [EK], which applies to strongly additive functions—additive functions f such that $f(mn) = f(m) + f(n)$ for all natural numbers m, n , and $|f(p)| \leq 1$ for all primes p —to immediately obtain the following statement.

Theorem 4.3 (Erdős-Kac). *Fix a modulus q and constants $A, B \in \mathbf{R}$. Then*

$$(4.9) \quad \lim_{x \rightarrow \infty} \frac{1}{x} \#\left\{n \leq x : A \leq \frac{\omega(n; q, a) - \frac{1}{\phi(q)} \log \log x}{\sqrt{\frac{1}{\phi(q)} \log \log x}} \leq B\right\} = \frac{1}{\sqrt{2\pi}} \int_A^B e^{-t^2/2} dt.$$

Hence $\omega(n; q, a)$ is also normally distributed.

FIGURE 5. Distribution of $\omega(n; q, a)$.

5. NUMERICAL TESTS

In this section, we discuss numerical observations regarding the behaviour of the summatory Liouville functions for arithmetic progressions, and also $\lambda(n; q, a_1, \dots, a_r)$ for mixed residue classes. We summarize our observations in Problem 5.1.

5.1. Numerical methods. All computations in this paper were done on Fortran95 and Python3. The largest computation that we could carry out was that of $L(x; 4, 1)$, which we verified to be positive for all $1 < x \leq 10^{11}$. Most of the other calculations were carried out up to 10^8 or 10^9 . Because of the limited amount of computer memory, we used an algorithm that calculates the parities in batches of size 10^8 . Also, to avoid the problem of factorizing large integers, we used multiplication to build up the parities of numbers up to x . This results in a significant increase in the speed without any type of parallel computation. The codes for the computations of various combinations were primarily written in Python, which allows to easily construct all the required combinations.

From the figures in the Introduction, and further numerical experiments, we pose the following problems (rather than conjectures, in light of Pólya and Turán):

Problem 5.1. (1) Let $a_1, \dots, a_{\varphi(q)}$ be the residue classes coprime to q , as in (3.22). Then is

$$(5.1) \quad \sum_{n \leq x} \lambda(n; q, a_1, \dots, a_{\varphi(q)}) \leq 0$$

for all x large enough?




FIGURE 6. Mixing residue classes.

(2) Let $\lambda(n; q, a_1, \dots, a_r)$ be complementary to a Dirichlet character mod q , as in (3.55). Then is

$$(5.2) \quad \sum_{n \leq x} \lambda(n; q, a_1, \dots, a_r) \geq 0$$

for all x large enough?

(3) What is the effect of mixing residue classes? (See 5.1.1.)

Also, one can ask in each case when the sign changes are eventually constant, if there are in fact no sign changes at all to the sums.

5.1.1. Mixing residue classes. As exemplified in the Figure 6 above, we notice that the addition of residue classes 2 and 3, which divide 6, affect the fluctuations in the sum in a nontrivial manner. In fact, while we know that $L(x; 6, 1)$ is asymptotically positive, it appears that $L(x; 6, 1, 2, 3)$ should change sign infinitely often.

5.2. Speculations on sieve parity. We close with a vague speculation regarding sieve parity, following the heuristic of Tao [T, 3.10.2] regarding the parity problem, in terms of the Liouville function. Let A be a set we would like to sieve for. Then to get a lower bound on $|A|$ in say, $[x, 2x]$ we set up the divisor sum lower bound

$$(5.3) \quad 1_A(n) \geq \sum_{d|n} c_d$$

where the divisors are concentrated in some sieve level $d \leq R$. Summing over $n \leq x$ we get the form

$$(5.4) \quad |A| \geq \sum_{d|n} c_d \frac{x}{d} + \dots$$

If instead we multiply by the nonnegative weight $1 + \lambda(n; q, a)$ and sum, we obtain

$$(5.5) \quad 2|A \setminus A_{a,q}| \geq \sum_{n \leq x} \sum_{d|n} c_d (1 + \lambda(n; q, a)) = \sum_{d|n} c_d \frac{x}{d} + L(x; q, a) + \dots$$

where $A_{a,q}$ denotes the set of $x \in A$ such that $x \equiv a \pmod{q}$. This suggests that one may be able to produce nontrivial lower bounds on the complement $|A \setminus A_{a,q}|$ by sieve theory, when $\varphi(q) > 2$.

In this light, we mention the result of [RW] who prove, under certain conditions, the existence of a product of three primes below $x^{1/3}$, each congruent to $a \pmod{q}$ where $q \leq x^{1/16}$. In particular, they observe that their use of a sieve is not blocked by sieve parity. More importantly, in a future work we hope to relate $\lambda(n; q, a)$ to the Rosser-Iwaniec sieve, in which the sieve $S(A, z)$ is bounded by sets defined by the sign of $\lambda(n)$.

REFERENCES

- [AGS] A. Balog, A. Granville and K. Soundararajan, Multiplicative functions in arithmetic progressions, *Ann. Math. Qué.* **37** (2013), no. 1, 3–30. MR3117735
- [BCC] P. Borwein, S. K. K. Choi and M. Coons, Completely multiplicative functions taking values in $\{-1, 1\}$, *Trans. Amer. Math. Soc.* **362** (2010), no. 12, 6279–6291. MR2678974
- [C] M. E. Changa, On the sums of multiplicative functions over numbers, all of whose divisors lie in a given arithmetic progression, *Izv. Math.* **69** (2005), no. 2, 423–438; translated from *Izv. Ross. Akad. Nauk Ser. Mat.* **69** (2005), no. 2, 205–220. MR2136261
- [EK] P. Erdős and M. Kac, The Gaussian law of errors in the theory of additive number theoretic functions, *Amer. J. Math.* **62** (1940), 738–742. MR0002374
- [H] C. B. Haselgrove, A disproof of a conjecture of Pólya, *Mathematika* **5** (1958), 141–145. MR0104638
- [HS] K. Halupczok and B. Suger, Partial sums of the Möbius function in arithmetic progressions assuming GRH, *Funct. Approx. Comment. Math.* **48** (2013), part 1, 61–90. MR3086961
- [K] A. A. Karatsuba, On a property of the set of prime numbers, *Russian Math. Surveys* **66** (2011), no. 2, 209–220; translated from *Uspekhi Mat. Nauk* **66** (2011), no. 2(398), 3–14. MR2847788
- [LOS] R. J. Lemke Oliver and K. Soundararajan, Unexpected biases in the distribution of consecutive primes, *Proc. Natl. Acad. Sci. USA* **113** (2016), no. 31, E4446–E4454. MR3624386
- [MR] K. Matomäki and M. Radziwiłł, Multiplicative functions in short intervals, *Ann. of Math.* (2) **183** (2016), no. 3, 1015–1056. MR3488742
- [MRT] M. J. Mossinghoff and T. S. Trudgian, Between the problems of Pólya and Turán, *J. Aust. Math. Soc.* **93** (2012), no. 1-2, 157–171. MR3062002
- [RW] O. Ramaré and A. Walker, Products of primes in arithmetic progressions: a footnote in parity breaking, preprint.
- [T] T. Tao, Structure and randomness. Pages from year one of a mathematical blog. American Mathematical Society, Providence, RI, 2008. xii+298 pp. ISBN: 978-0-8218-4695-7 MR2459552
- [Y] L. Ye., Bounding sums of the Möbius function over arithmetic progressions, arXiv:1406.7326v1.

E-mail address: snehal.s@iiserpune.ac.in

INDIAN INSTITUTE FOR SCIENCE EDUCATION AND RESEARCH (IISER) PUNE, DR HOMI BHABHA
ROAD, PASHAN, PUNE, 411008 INDIA

E-mail address: tiananwong@iiserpune.ac.in

INDIAN INSTITUTE FOR SCIENCE EDUCATION AND RESEARCH (IISER) PUNE, DR HOMI BHABHA
ROAD, PASHAN, PUNE, 411008 INDIA